

Short Communication

## Correlation and path analysis studies in parthenocarpic cucumber (*Cucumis sativus* L.)

Palvi Mehta, Parveen Sharma\*, Manpreet Kaur, Shilpa, Akhilesh Sharma and Neelam Bhardwaj<sup>1</sup>

Received: March 2022/ Accepted: July 2022

Cucumber (*Cucumis sativus* L.) is the most important member of cucurbitaceae family due to its economic importance. It helps to cure kidney infection, jaundice, stomach problems and indigestion. It is believed to be originated in India and the wild relative *Cucumis sativus* var. *hardwickii* is the progenitor of the cultivated cucumber (Choudhary et al. 2015; Sharma 2017). It is most suitable vegetable for protected conditions due to year round production and higher demand in market. Due to its origin in India, it has accumulated most of the genetic variability which helps to improve various characters through selection. Germplasm purity is difficult to maintain due to higher cross pollination in cucumber. It depicts wide range of variability with no uniformity in characters like fruit size, shape, color and yield among existing germplasm (Sharma et al. 2017; Kaur and Sharma 2022). Parthenocarpic genotypes are widely grown under protected conditions which bear female flowers in every node and develop seedless fruits. Presence of genetic variability in germplasm helps in crop improvement through selection. Therefore, genetic restructuring of cucumber germplasm is the first step to identify the potential genotypes for use in breeding programme. Most of economic traits are quantitative in nature, which are influenced by the environment and their effective selection relies on the nature of genetic and non-genetic variation. It will help in partitioning the overall variability into heritability and non-heritable components. Understanding of interrelationships among various characters is useful because selection of one character may affect the performance of other characters, which helps in identification of components

of complex characters such as yield. Determination of correlation does not give a correct depiction of the direct influence of each of the components traits towards the yield. Path coefficient helps in partitioning the correlation coefficient into direct and indirect effects and helps in identification of characters which are useful selection criteria to improve fruit yield. Thus, present investigation was carried out to study the character association and direct & indirect effects of independent characters on yield.

The experiment was conducted at Department of Vegetable Science and Floriculture, College of Agriculture, CSKHPKV, Palampur during spring-summer 2018 under naturally ventilated polyhouse. The experimental farm is situated at 32°6' N latitude, 76°3' E longitude and at an elevation of 1,290.8 m above mean sea level. The experimental material consists of twelve genotypes of parthenocarpic cucumber along with two checks were evaluated in Randomized Block Design (RBD) with three replications in modified naturally ventilated poly-house. Ten plants in each replication were planted at spacing of 70×30 cm within row to row and plant to plant, respectively to determine components of variability, association of various traits with yield and their direct and indirect effects for successful selection for crop improvement. Data were recorded on randomly taken ten plants of each genotype in each replication on growth and yield contributing characters viz., days to anthesis of first female flower, nodal position of first female flower, number of female flowers per node, days taken to first fruit harvest, fruit length (cm), fruit girth (cm), fruit weight (g), number of fruits per plant, marketable yield per plant (kg), harvest duration (days), internodal length (cm), vine length (m), total soluble solids (°Brix), incidence of powdery mildew disease (%) and incidence of downy mildew disease (%). Correlation coefficients analysis were carried as per the method of Al-Jibouri et al (1958) while, path coefficient analysis

---

Department of Vegetable Science and Floriculture, CSK Himachal Pradesh Agricultural University, Palampur, HP  
<sup>1</sup>Rice &Wheat Research Centre, Malan, CSK Himachal Pradesh Agricultural University, Palampur, HP  
\*Corresponding author, E-mail: parveens012@gmail.com



**Table 2:** Estimation of direct and indirect effects of different traits on marketable yield per plant at phenotypic (P) and genotypic (G) levels in parthenocarpic cucumber genotypes

Traits		Days to anthesis of first female flower	Nodal position of first female flower	Number of female flowers per node	Days taken to first fruit harvest	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)	Number of fruits per plant	Harvest duration (days)	Internodal length (cm)	Vine length (m)	Total soluble solids ( $^{\circ}$ Brix)	r
Days to anthesis of first female flower	P	<b>-0.018</b>	0.008	0.000	0.015	0.006	-0.004	-0.316	-0.297	-0.007	0.004	-0.004	0.009	-0.603**
	G	<b>-0.007</b>	0.012	-0.001	0.011	0.013	-0.035	-0.456	-0.412	-0.003	-0.017	-0.035	0.015	-0.915**
Nodal position of first female flower	P	-0.004	<b>0.031</b>	0.001	0.007	0.006	-0.002	-0.165	-0.211	-0.003	0.004	-0.003	-0.006	-0.345*
	G	-0.005	<b>0.017</b>	0.008	0.006	0.016	-0.018	-0.266	-0.277	-0.001	-0.012	-0.019	-0.006	-0.558**
Number of female flowers per node	P	0.000	0.005	<b>0.003</b>	0.000	0.001	-0.002	0.016	0.010	-0.003	0.001	-0.002	-0.001	0.029
	G	0.000	0.006	<b>0.022</b>	0.001	0.000	-0.004	0.023	-0.035	-0.003	-0.004	-0.012	-0.001	-0.005
Days taken to first fruit harvest	P	-0.010	0.008	0.000	<b>0.027</b>	0.006	-0.008	-0.389	-0.409	-0.010	0.005	-0.003	0.009	-0.773**
	G	-0.005	0.007	0.002	<b>0.014</b>	0.015	-0.055	-0.461	-0.460	-0.006	-0.015	-0.027	0.017	-0.976**
Fruit length (cm)	P	0.008	-0.014	0.000	-0.012	<b>-0.013</b>	0.007	0.284	0.259	0.007	-0.005	0.004	-0.002	0.524**
	G	0.005	-0.016	0.000	-0.012	<b>-0.017</b>	0.023	0.394	0.394	0.006	0.016	0.030	-0.001	0.823**
Fruit girth (cm)	P	0.003	-0.003	0.000	-0.011	-0.005	<b>0.020</b>	0.283	0.347	0.004	-0.005	0.004	-0.010	0.628**
	G	0.004	-0.005	-0.001	-0.013	-0.007	<b>0.060</b>	0.402	0.478	0.004	0.015	0.028	-0.008	0.957**
Fruit weight (g)	P	0.011	-0.010	0.000	-0.021	-0.008	0.011	<b>0.505</b>	0.481	0.007	-0.006	0.005	-0.012	0.962**
	G	0.007	-0.010	0.001	-0.014	-0.015	0.053	<b>0.457</b>	0.464	0.006	0.016	0.029	-0.012	0.983**
Number of fruits per plant	P	0.009	-0.012	0.000	-0.020	-0.006	0.012	0.441	<b>0.550</b>	0.008	-0.006	0.004	-0.011	0.970**
	G	0.006	-0.009	-0.002	-0.013	-0.014	0.059	0.436	<b>0.487</b>	0.005	0.017	0.031	-0.012	0.991**
Harvest duration (days)	P	0.007	-0.005	-0.001	-0.014	-0.005	0.004	0.182	0.219	<b>0.019</b>	-0.003	0.001	-0.007	0.398*
	G	0.003	-0.003	-0.008	-0.010	-0.013	0.028	0.332	0.322	<b>0.008</b>	0.018	0.019	-0.011	0.685**
Internodal length (cm)	P	-0.007	0.011	0.000	0.014	0.006	-0.010	-0.319	-0.339	-0.006	<b>0.010</b>	-0.006	0.009	-0.636**
	G	-0.006	0.011	0.005	0.012	0.016	-0.049	-0.414	-0.472	-0.008	<b>-0.018</b>	-0.030	0.010	-0.946**
Vine length (m)	P	0.009	-0.011	-0.001	-0.010	-0.006	0.010	0.256	0.261	0.003	-0.007	<b>0.009</b>	-0.009	0.504**
	G	0.006	-0.009	-0.007	-0.010	-0.014	0.045	0.361	0.405	0.004	0.015	<b>0.037</b>	-0.012	0.822**
Total soluble solids ( $^{\circ}$ Brix)	P	-0.006	-0.006	0.000	0.009	0.001	-0.007	-0.210	-0.211	-0.005	0.003	-0.003	<b>0.028</b>	-0.407*
	G	-0.004	-0.004	-0.001	0.009	0.001	-0.019	-0.219	-0.240	-0.003	-0.007	-0.018	<b>0.024</b>	-0.482**

also reported direct and positive effects of number of fruits per plant, fruit weight, number of female flower per node and days taken to first fruit harvest, nodal position of first female flower, harvest duration, fruit girth and total soluble solids. However, marketable yield per plant was also significantly increased by maximum positive indirect effects of fruit girth via number of fruits per plant followed by fruit weight via number of fruits per plant. As number of fruits per plant, fruit weight and fruit girth have highest selection index thereby more emphasis needed to be given for inclusion of these traits in parthenocarpic cucumber improvement programme. In the present investigation, number of fruits per plant has exhibited highly significant positive association with marketable yield per plant followed by fruit weight, fruit girth, fruit length, vine length and harvest duration. The path coefficient analysis revealed that the number of fruits per plant and fruit weight had direct positive phenotypic and genotypic effect on yield. These findings showed that direct selection on the basis of number of fruits per plant and fruit weight will be rewarding for crop improvement in parthenocarpic cucumber.

## References

- Al-Jibouri HA, Miller PA and Robinson HF (1958) Genotypic and environmental variance and co-variance in upland cotton crops of inter-specific origin. *Agron J* 50: 633-636.
- Choudhary H, Singh DK and Damke SR (2015) Genetic variability in *Cucumis sativus* var. *hardwickii*: key to cucumber improvement. *Int J Basic Appl Agric Res* 13: 340-343.
- Dewey JR and Lu KH (1959) Correlation and path analysis of components of crested wheat grass seed production. *Agron J* 51: 515-518.
- Hasan R, Hossain MK, Alum N, Bashar A, Islam S and Tarafder MJA (2015) Genetic divergence in commercial cucumber (*Cucumis sativus* L.) genotypes. *Bangladesh J Bot* 44: 201-207.
- Kaur M and Sharma P (2022) Recent advances in cucumber (*Cucumis sativus* L.). *J Hortic Sci Biotech* 97: 3-23.
- Kumar P, Khapte PS, Saxena A and Kumar (2019) Evaluation of gynocercous cucumber (*Cucumis sativus*) hybrids for early-summer greenhouse production in Western Indian arid plains. *Indian J Agric Sci* 89: 545-550.
- Monge PJE, Chacon PK and Loria CM (2021) Selection criteria for yield in cucumber (*Cucumis sativus*) grown under greenhouse conditions during the dry season. *UNED Res J* 13: e3373.

- Pal S (2016) Genetic studies for yield and quality traits in cucumber (*Cucumis sativus* L.). M.Sc. Thesis, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, HP.
- Sharma S (2017) Genetic divergence studies in cucumber (*Cucumis sativus* L.). M.Sc. Thesis, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, HP.
- Sharma S, Kumar R and Sharma HR (2017) Studies on Variability, Heritability and Genetic Gain in Cucumber (*Cucumis sativus* L.). I J Ecol 44: 829-833.
- Singh SS, Yadav GC and Kathayat K (2018) Study of genetic variability in cucumber. J Hill Agric 9: 39-43.
- Singh Y, Safiullah, Verma A, Sharma S and Sekhon BS (2017) Genetic evaluation of cucumber (*Cucumis sativus* L.) genotypes for yield and its contributing traits under mid-hill conditions of Himachal Pradesh, India. Environ & Ecol 35: 3621-26.
- Sonia S, Naveen K, Chandel KS and Parveen S (2011) Determination of genetic variation for morphological and yield traits in bell pepper (*Capsicum annuum* var. *grossum*). Indian J Agri Sci 81(7): 590-594.
- Verma S (2003) Genetic variability and correlation studies in cucumber (*Cucumis sativus* L.). M.Sc. Thesis, Dr. YS Parmar University of Horticulture and Forestry, Nauni, Solan, HP.